

Immunization schedule

	Birth	2 months	4 months	6 months	12 to 15 months	18 to 23 months	4 to 6 years	12 years	14 to 18 years	19 to 49 years	50 to 64 years	65+ years
Hepatitis B	X											
Rotavirus		X	X	X								
DTaP-HepB-IPV ¹		X	X	X								
H. influenza type B (Hib)		X	X	X	X							
Pneumococcal conjugate (PCV7)		X	X	X	X							
DTaP					X		X					
IPV							X					
Measles, mumps, rubella (MMR)					X		X					
Varicella (chickenpox)					X		X	X ²				
Hepatitis A					X	X						
Influenza				Every year						Every year ³	Every year	
Tetanus, diphtheria, pertussis (Tdap)								X	Every 10 years			
									Substitute 1 dose of Tdap for Td			
Meningococcal conjugate (MCV4)								X	X ⁴			
Human papillomavirus (HPV)								X ⁵	X ^{4,5}			
Pneumococcal polysaccharide										X ³	X ³	X
Shingles (zoster)												X ⁶

¹ Combination vaccine includes diphtheria, tetanus, pertussis, hepatitis B and polio.

² All people who have not had chickenpox disease should receive two doses of vaccine.

³ Recommended for people with chronic health problems.

⁴ If not previously administered.

⁵ Females age 9 through 26 only: three-dose series, second dose two months after first dose, third dose four months after second dose.

⁶ Recommended for people age 60 years or older.

Revised: 7/2/08